

Nested Padé approximants as a tool for multivariate finite element problems with porous media

SAPEM 2014, Stockholm, December 16, 2014

- Romain Rumpler* – CREO Dynamics AB, Linköping
Peter Göransson – KTH, Stockholm
Henry Rice – Trinity College Dublin





Motivations and objectives

- Establish a Padé-based approach for fast parametric sweeps of general problems
- Extend the method to multivariate problems
- Account for non trivial parametric analyses
- Potential applications:
 - ★ Optimization problems
 - ★ Response for stochastic systems
 - ★ ...



ROYAL INSTITUTE OF TECHNOLOGY



Motivations

An extension to multivariate expansions?

Some approaches from the literature

The nested Padé approximants

Test case: acoustic absorption

Bivariate Padé reconstruction

Conclusion

A few approaches of potential interest...

- Homogeneous multivariate Padé approximants [Cuyt, 1983,1985,1997]
 - ✓ Efficient recursive algorithms developed
 - ✓ Convergence proofs
 - ✗ Rapid increase of numerator and denominator polynomial degrees
 - ✗ Singularities potentially appearing in the convergence domain



A few approaches of potential interest...

- Homogeneous multivariate Padé approximants [Cuyt, 1983,1985,1997]
 - ✓ Efficient recursive algorithms developed
 - ✓ Convergence proofs
 - ✗ Rapid increase of numerator and denominator polynomial degrees
 - ✗ Singularities potentially appearing in the convergence domain
- Composition of univariate Padé approximants [Chaffy, 1988]
 - ✓ Based on the calculation of univariate Padé approximants
 - ✓ Convergence proofs and convenient implementation
 - ✗ Recursive increase of the polynomial degrees of the multivariate approximant

A few approaches of potential interest...

- Homogeneous multivariate Padé approximants [Cuyt, 1983,1985,1997]
 - ✓ Efficient recursive algorithms developed
 - ✓ Convergence proofs
 - ✗ Rapid increase of numerator and denominator polynomial degrees
 - ✗ Singularities potentially appearing in the convergence domain
- Composition of univariate Padé approximants [Chaffy, 1988]
 - ✓ Based on the calculation of univariate Padé approximants
 - ✓ Convergence proofs and convenient implementation
 - ✗ Recursive increase of the polynomial degrees of the multivariate approximant
- Nested Padé approximants [Guillaume, 1997]
 - ✓ Univariate Padé approximants, convergence proofs and straightforward implementation
 - ✓ As many final coefficients as the initial multivariate Taylor expansion
 - ✗ Performance convergence/cost?



Motivations

An extension to multivariate expansions?

Some approaches from the literature

The nested Padé approximants

Test case: acoustic absorption

Bivariate Padé reconstruction

Conclusion

The nested Padé approximants: main steps, bivariate case (1/3)

- General form of the problems to solve: $\mathbf{Z}(x, y)\mathbf{u}(x, y) = \mathbf{f}(x, y)$



The nested Padé approximants: main steps, bivariate case (1/3)



- General form of the problems to solve: $\mathbf{Z}(x, y)\mathbf{u}(x, y) = \mathbf{f}(x, y)$

- Expansion as a double power series:

$$\star U(x_0 + \Delta x, y_0 + \Delta y) \approx \sum_{i=0}^{N_i} \sum_{j=0}^{N_j} r_{ij} \Delta x^i \Delta y^j, \text{ with } r_{ij} = \frac{U^{(i)(j)}(x_0, y_0)}{i!j!}$$

★ Involves $(N_i + 1) * (N_j + 1)$ coefficients r_{ij}

★ $U^{(i)(j)}(x_0, y_0)$ recursively calculated via a Liebnitz formula (multiple RHS)



The nested Padé approximants: main steps, bivariate case (1/3)

- General form of the problems to solve: $\mathbf{Z}(x, y)\mathbf{u}(x, y) = \mathbf{f}(x, y)$

- Expansion as a double power series:

$$\star U(x_0 + \Delta x, y_0 + \Delta y) \approx \sum_{i=0}^{N_i} \sum_{j=0}^{N_j} r_{ij} \Delta x^i \Delta y^j, \text{ with } r_{ij} = \frac{U^{(i)(j)}(x_0, y_0)}{i!j!}$$

★ Involves $(N_i + 1) * (N_j + 1)$ coefficients r_{ij}

★ $U^{(i)(j)}(x_0, y_0)$ recursively calculated via a Liebnitz formula (multiple RHS)

- Rewriting as an intermediate (Δy -parametrized) Padé approximant:

$$\star U_{\Delta y}(\Delta x) = \frac{P(\Delta x, \Delta y)}{Q(\Delta x, \Delta y)} = \frac{\sum_{i=0}^{m_i} s_i^{N_j}(\Delta y) \Delta x^i}{1 + \sum_{i=1}^{n_i} s_{m_i+i}^{N_j}(\Delta y) \Delta x^i}, \text{ with } m_i + n_i = N_i$$



The nested Padé approximants: main steps, bivariate case (1/3)

- General form of the problems to solve: $\mathbf{Z}(x, y)\mathbf{u}(x, y) = \mathbf{f}(x, y)$

- Expansion as a double power series:

$$\star U(x_0 + \Delta x, y_0 + \Delta y) \approx \sum_{i=0}^{N_i} \sum_{j=0}^{N_j} r_{ij} \Delta x^i \Delta y^j, \text{ with } r_{ij} = \frac{U^{(i)(j)}(x_0, y_0)}{i!j!}$$

★ Involves $(N_i + 1) * (N_j + 1)$ coefficients r_{ij}

★ $U^{(i)(j)}(x_0, y_0)$ recursively calculated via a Liebnitz formula (multiple RHS)

- Rewriting as an intermediate (Δy -parametrized) Padé approximant:

$$\star U_{\Delta y}(\Delta x) = \frac{P(\Delta x, \Delta y)}{Q(\Delta x, \Delta y)} = \frac{\sum_{i=0}^{m_i} s_i^{N_j}(\Delta y) \Delta x^i}{1 + \sum_{i=1}^{n_i} s_{m_i+i}^{N_j}(\Delta y) \Delta x^i}, \text{ with } m_i + n_i = N_i$$

- Intermediate Padé coefficients:

$$\star s_i^{N_j}(\Delta y) = \sum_{j=0}^{N_j} s_{ij} \Delta y^j \implies s_{ij}?$$



The nested Padé approximants: main steps, bivariate case (1/3)

- General form of the problems to solve: $\mathbf{Z}(x, y)\mathbf{u}(x, y) = \mathbf{f}(x, y)$

- Expansion as a double power series:

$$\star U(x_0 + \Delta x, y_0 + \Delta y) \approx \sum_{i=0}^{N_i} \sum_{j=0}^{N_j} r_{ij} \Delta x^i \Delta y^j, \text{ with } r_{ij} = \frac{U^{(i)(j)}(x_0, y_0)}{i! j!}$$

★ Involves $(N_i + 1) * (N_j + 1)$ coefficients r_{ij}

★ $U^{(i)(j)}(x_0, y_0)$ recursively calculated via a Liebnitz formula (multiple RHS)

- Rewriting as an intermediate (Δy -parametrized) Padé approximant:

$$\star U_{\Delta y}(\Delta x) = \frac{P(\Delta x, \Delta y)}{Q(\Delta x, \Delta y)} = \frac{\sum_{i=0}^{m_i} s_i^{N_j}(\Delta y) \Delta x^i}{1 + \sum_{i=1}^{n_i} s_{m_i+i}^{N_j}(\Delta y) \Delta x^i}, \text{ with } m_i + n_i = N_i$$

- Intermediate Padé coefficients:

$$\star s_i^{N_j}(\Delta y) = \sum_{j=0}^{N_j} s_{ij} \Delta y^j \implies s_{ij} ? \implies \begin{cases} \mathbf{H}_0 \mathbf{S}_0 = \mathbf{C}_0 \\ \mathbf{H}_0 \mathbf{S}_j = \mathbf{C}_j - \sum_{k=1}^j \mathbf{H}_k \mathbf{S}_{j-k}, \quad j = 1 \dots N_j \end{cases}$$

The nested Padé approximants: main steps, bivariate case (2/3)



- Rewriting as a nested Padé approximant:

$$\star U_{\Delta y}(\Delta x) = \frac{\sum_{i=0}^{m_i} s_i^{N_j}(\Delta y) \Delta x^i}{1 + \sum_{i=1}^{n_i} s_{m_i+i}^{N_j}(\Delta y) \Delta x^i}, \text{ with } s_i^{N_j}(\Delta y) = \sum_{j=0}^{N_j} s_{ij} \Delta y^j$$

$$\star s_i^{N_j}(\Delta y) \text{ with Padé} \implies s_i^{N_j}(\Delta y) = \frac{\sum_{j=0}^{m_j} t_{ij} \Delta y^j}{1 + \sum_{j=1}^{n_j} t_{i(m_j+j)} \Delta y^j}, \text{ with } m_j + n_j = N_j$$

The nested Padé approximants: main steps, bivariate case (2/3)



- Rewriting as a nested Padé approximant:

$$\star U_{\Delta y}(\Delta x) = \frac{\sum_{i=0}^{m_i} s_i^{N_j}(\Delta y) \Delta x^i}{1 + \sum_{i=1}^{n_i} s_{m_i+i}^{N_j}(\Delta y) \Delta x^i}, \text{ with } s_i^{N_j}(\Delta y) = \sum_{j=0}^{N_j} s_{ij} \Delta y^j$$

$$\star s_i^{N_j}(\Delta y) \text{ with Padé} \implies s_i^{N_j}(\Delta y) = \frac{\sum_{j=0}^{m_j} t_{ij} \Delta y^j}{1 + \sum_{j=1}^{n_j} t_{i(m_j+j)} \Delta y^j}, \text{ with } m_j + n_j = N_j$$

May be rewritten as a system of linear equations:

$$[\mathbf{A}_i] \begin{bmatrix} t_{i(m_j+1)} \\ \vdots \\ t_{i(N_j)} \\ t_{i0} \\ \vdots \\ t_{i(m_j)} \end{bmatrix} = \begin{bmatrix} s_{i0} \\ \vdots \\ s_{i(n_j-1)} \\ s_{i(n_j)} \\ \vdots \\ s_{i(N_j)} \end{bmatrix}, \text{ for } i = 1, \dots, N_i + 1$$

The nested Padé approximants: main steps, bivariate case (3/3)



ROYAL INSTITUTE
OF TECHNOLOGY

- Nested Padé solution expansion around (x_0, y_0) , for $\mathbf{Z}(x_0, y_0)\mathbf{u}(x_0, y_0) = \mathbf{f}(x_0, y_0)$?

The nested Padé approximants: main steps, bivariate case (3/3)



- Nested Padé solution expansion around (x_0, y_0) , for $\mathbf{Z}(x_0, y_0)\mathbf{u}(x_0, y_0) = \mathbf{f}(x_0, y_0)$?
- Steps of the procedure at (x_0, y_0) :
 - ▶ Compute **matrix decomposition** of $\mathbf{Z}(x_0, y_0)$

The nested Padé approximants: main steps, bivariate case (3/3)



- Nested Padé solution expansion around (x_0, y_0) , for $\mathbf{Z}(x_0, y_0)\mathbf{u}(x_0, y_0) = \mathbf{f}(x_0, y_0)$?
- Steps of the procedure at (x_0, y_0) :
 - ▶ Compute **matrix decomposition** of $\mathbf{Z}(x_0, y_0)$
 - ▶ **Solve** for $\mathbf{u}(x_0, y_0)$

The nested Padé approximants: main steps, bivariate case (3/3)



- Nested Padé solution expansion around (x_0, y_0) , for $\mathbf{Z}(x_0, y_0)\mathbf{u}(x_0, y_0) = \mathbf{f}(x_0, y_0)$?
- Steps of the procedure at (x_0, y_0) :
 - ▶ Compute **matrix decomposition** of $\mathbf{Z}(x_0, y_0)$
 - ▶ **Solve** for $\mathbf{u}(x_0, y_0)$
 - ▶ **Solve** recursively for the first partial derivatives of $\mathbf{u}^{(i)(j)}(x_0, y_0)$,
 $[(N_i + 1) * (N_j + 1) - 1]$ multiple RHS full solutions

The nested Padé approximants: main steps, bivariate case (3/3)



- Nested Padé solution expansion around (x_0, y_0) , for $\mathbf{Z}(x_0, y_0)\mathbf{u}(x_0, y_0) = \mathbf{f}(x_0, y_0)$?
- Steps of the procedure at (x_0, y_0) :
 - ▶ Compute **matrix decomposition** of $\mathbf{Z}(x_0, y_0)$
 - ▶ **Solve** for $\mathbf{u}(x_0, y_0)$
 - ▶ **Solve** recursively for the first partial derivatives of $\mathbf{u}^{(i)(j)}(x_0, y_0)$,
 $[(N_i + 1) * (N_j + 1) - 1]$ multiple RHS full solutions
 - ▶ **Solve** recursively for the intermediate Padé coefficients at dofs of interest
 $(N_j + 1)$ systems of size n_j

The nested Padé approximants: main steps, bivariate case (3/3)



- Nested Padé solution expansion around (x_0, y_0) , for $\mathbf{Z}(x_0, y_0)\mathbf{u}(x_0, y_0) = \mathbf{f}(x_0, y_0)$?
- Steps of the procedure at (x_0, y_0) :
 - ▶ Compute **matrix decomposition** of $\mathbf{Z}(x_0, y_0)$
 - ▶ **Solve** for $\mathbf{u}(x_0, y_0)$
 - ▶ **Solve** recursively for the first partial derivatives of $\mathbf{u}^{(i)(j)}(x_0, y_0)$,
 $[(N_i + 1) * (N_j + 1) - 1]$ multiple RHS full solutions
 - ▶ **Solve** recursively for the intermediate Padé coefficients at dofs of interest
 $(N_j + 1)$ systems of size n_j
 - ▶ **Solve** for the nested Padé coefficients
 $(N_j + 1)$ systems of size n_j

The nested Padé approximants: main steps, bivariate case (3/3)



- Nested Padé solution expansion around (x_0, y_0) , for $\mathbf{Z}(x_0, y_0)\mathbf{u}(x_0, y_0) = \mathbf{f}(x_0, y_0)$?
- Steps of the procedure at (x_0, y_0) :
 - ▶ Compute **matrix decomposition** of $\mathbf{Z}(x_0, y_0)$
 - ▶ **Solve** for $\mathbf{u}(x_0, y_0)$
 - ▶ **Solve** recursively for the first partial derivatives of $\mathbf{u}^{(i)(j)}(x_0, y_0)$, $[(N_i + 1) * (N_j + 1) - 1]$ multiple RHS full solutions
 - ▶ **Solve** recursively for the intermediate Padé coefficients at dofs of interest ($N_j + 1$ systems of size n_j)
 - ▶ **Solve** for the nested Padé coefficients ($N_j + 1$ systems of size n_j)
 - ▶ **Rebuild solution** around (x_0, y_0) : $\mathbf{u}(x_0 + \Delta x, y_0 + \Delta y)$ (simple algebraic evaluation)

Motivations

An extension to multivariate expansions?

Some approaches from the literature

The nested Padé approximants


Test case: acoustic absorption

Bivariate Padé reconstruction

Conclusion


Test case: equivalent fluid porous material

- Absorption problem:




Acoustic impedance (Real part), $(f_0, \sigma_0) = (2500 \text{ Hz}, 25000 \text{ N.s/m}^4)$

- Reference solution




- 1-point bivariate reconstruction






Acoustic impedance (Real part), $(f_0, \sigma_0) = (2500 \text{ Hz}, 25000 \text{ N.s/m}^4)$


- Reference solution




- 1-point bivariate reconstruction



- Relative error ($N_f = N_\sigma = 9$)




- Relative error ($N_f = N_\sigma = 9$)




Absorption coefficient, $(f_0, \sigma_0) = (2500 \text{ Hz}, 25000 \text{ N.s/m}^4)$


- Reference/reconstructed solution



- Absorption curves



- Relative error ($N_f = N_\sigma = 9$)





Motivations

An extension to multivariate expansions?

Bivariate Padé reconstruction

Reconstruction procedure


3D elasto-poro-acoustic problem

Conclusion

Reconstruction steps over multiple intervals



Steps 1–3





Reconstruction steps over multiple intervals



ROYAL INSTITUTE OF TECHNOLOGY


Steps 1–3





Reconstruction steps over multiple intervals

Steps 1–3



Subsequent steps & further details:

[Rumpler R., Göransson P., Deü J.-F.], *A finite element approach combining a reduced-order system, Padé approximants, and an adaptive frequency windowing for fast multi-frequency solution of poro-acoustic problems*, IJNME, 97(10):1097-0207, 2014.

[Rumpler R., Göransson P., Rice H.J.], *An adaptive strategy for the bivariate solution of finite element problems using multivariate nested Padé approximants*, IJNME, 100:689–710, 2014.



Motivations

An extension to multivariate expansions?

Bivariate Padé reconstruction


Reconstruction procedure

3D elasto-poro-acoustic problem

Conclusion


3D elasto-poro-acoustic problem

- Absorption problem:



3D elasto-poro-acoustic problem

- Absorption problem:




- Delany-Bazley-Miki empirical model, equivalent fluid
- Dependence on **two parameters**: frequency f and flow resistivity σ
- Density** and **bulk modulus**: $\tilde{\rho}_{eq}(f, \sigma)$, $\tilde{K}_{eq}(f, \sigma)$
- Form of the problem to be solved: $\mathbf{Q}_p(f, \sigma) \mathbf{P}(f, \sigma) = \mathbf{V}$

$$\left[\begin{array}{c|c} \mathbf{K}_s - 4\pi^2 f^2 \mathbf{M}_s & -\mathbf{H}_{sa} \\ \hline -\mathbf{H}_{sa}^T & \frac{1}{4\pi^2 f^2 \rho_0} \mathbf{K}_a + \frac{1}{4\pi^2 f^2 \tilde{\rho}_{eq}(f, \sigma)} \mathbf{K}_p - \\ & \frac{1}{K_0} \mathbf{M}_a - \frac{1}{\tilde{K}_{eq}(f, \sigma)} \mathbf{M}_p \end{array} \right] \begin{bmatrix} \mathbf{U}(f, \sigma) \\ \mathbf{P}(f, \sigma) \end{bmatrix} = \begin{bmatrix} \mathbf{F} \\ \mathbf{0} \end{bmatrix}$$




Sound pressure level surface response

- Reference solution




- Padé bivariate reconstruction




Sound pressure level surface response


- Reference solution




- Padé bivariate reconstruction



- Relative error ($N_f = N_\sigma = 7$)




- Residue norm ($N_f = N_\sigma = 7$)




CPU time comparison – Padé reconstruction



- CPU reference vs. Nested Padé



- Speedup factor



Conclusion



Main conclusions

- **Good potential** using Padé-based interpolatory approaches for univariate and multivariate problems:
 - ▶ Good convergence properties on univariate and bivariate test cases
 - ▶ Based on univariate Padé approximants
 - ▶ Suited for an efficient implementation
 - ▶ Particularly interesting for partial field reconstruction
- Provides a piece-wise analytical approximate solution
- A numerical tool with several potential engineering applications
- An adaptive reconstruction – no a priori knowledge of the solution required

Conclusion



Main conclusions

- **Good potential** using Padé-based interpolatory approaches for univariate and multivariate problems:
 - ▶ Good convergence properties on univariate and bivariate test cases
 - ▶ Based on univariate Padé approximants
 - ▶ Suited for an efficient implementation
 - ▶ Particularly interesting for partial field reconstruction
- Provides a piece-wise analytical approximate solution
- A numerical tool with several potential engineering applications
- An adaptive reconstruction – no a priori knowledge of the solution required

THANK YOU FOR YOUR ATTENTION!

